



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SCIENCE

FRIDAY, JULY 22, 1921

TYPES OF VALENCE

CONTENTS

<i>Types of Valence:</i> DR. IRVING LANGMUIR.....	59
<i>Professor H. Bruchmann:</i> PROFESSOR DOUGLAS HOUGHTON CAMPBELL	67
<i>Scientific Events:—</i>	
<i>Field Work of the Smithsonian Institution; The Medical School of the University of Virginia; The Science Club of the University of Texas; The Rochester Meeting of the Optical Society of America; American Engineers in Europe</i>	68
<i>Scientific Notes and News</i>	71
<i>University and Educational News</i>	72
<i>Discussion and Correspondence:—</i>	
<i>The Geographic Distribution of Hybrids:</i> DR. M. L. FERNALD. <i>Albinism in the Black Bear:</i> PAUL C. STANDLEY. <i>Bechhold's "Capillary Phenomenon" in Agriculture:</i> JEROME ALEXANDER	73
<i>Quotations:—</i>	
<i>The Royal Institution.....</i>	74
<i>Scientific Books:—</i>	
<i>Evermann and Clark on Lake Maxinkuckee:</i> T. L. HANKINSON.....	75
<i>Special Articles:—</i>	
<i>An Explanation of Liesegang's Rings:</i> DR. HUGH MCGUIGAN	78
<i>The North Carolina Academy of Science:</i> C. S. BRIMLEY	80

FROM the theory proposed by G. N. Lewis in 1916 and subsequently extended by the writer, it is clear¹ that the term valence has been used in the past to cover what we may now recognize as three distinct types of valence, viz.:

1. Positive valence: the number of electrons an atom can give up.
2. Negative valence: the number of electrons an atom can take up.
3. Covalence: the number of pairs of electrons an atom can share with its neighbors.

It was shown that these fundamental conceptions of valence as well as the actual numerical values of each type of valence for most of the elements could be derived from a few postulates regarding the structure of atoms. The following method of deriving these relationships, however, is not only much simpler than that previously given by the writer, but throws a new light on the relationships between the different types of valence.

We will take for granted the Rutherford type of atom, which consists of a positive nucleus surrounded by a number of electrons equal to the atomic number of the atom. We will also assume that Coulomb's law applies to the forces between the charged particles in the atom, but at the same time will recognize the existence of repulsive forces in atoms which prevent the electrons from falling into the nucleus. For the present purpose, however, it is immaterial whether the repulsive force is a dynamic force (centrifugal force) such as that assumed by Bohr, or is a static force as postulated by G. N. Lewis, J. J. Thomson or recently by the writer.²

We shall need to make only 3 postulates in

¹ Langmuir, *Jour. Amer. Chem. Soc.*, 41, 926 (1919), and *Jour. Ind. Eng. Chem.*, 12, 386 (1920).

² SCIENCE, 53, 290, Mar. 25, 1921.